

Potential Complementarity of High-Flavanol Cocoa Powder and Spirulina for Health Protection

Mark F. McCarty, Jorge Barroso-Aranda, and Francisco Contreras,
Oasis of Hope Hospital, Tijuana, Mexico

Abstract

Recent studies show that ingestion of flavanol-rich cocoa powder provokes increased endothelial production of nitric oxide - an effect likely mediated by epicatechin – and thus may have considerable potential for promoting vascular health. The Kuna Indians of Panama, who regularly consume large amounts of flavanol-rich cocoa, are virtually free of hypertension and stroke, even though they salt their food. Of potentially complementary merit is the cyanobacterium spirulina, which has been used as a food in certain cultures. Spirulina is exceptionally rich in phycocyanobilin (PCB), which recently has been shown to act as a potent inhibitor of NADPH oxidase; this effect likely rationalizes the broad range of anti-inflammatory, cytoprotective, and anti-atherosclerotic effects which orally administered spirulina has achieved in rodent studies. In light of the central pathogenic role which NADPH oxidase-derived oxidant stress plays in a vast range of disorders, spirulina or PCB-enriched spirulina extracts may have remarkable potential for preserving and restoring health. Joint administration of flavanol-rich cocoa powder and spirulina may have particular merit, inasmuch as cocoa can mask the somewhat disagreeable flavor and odor of spirulina, whereas the antioxidant impact of spirulina could be expected to amplify the bioactivity of the nitric oxide evoked by cocoa flavanols in inflamed endothelium. Moreover, there is reason to suspect that, by optimizing cerebrovascular perfusion while quelling cerebral oxidant stress, cocoa powder and spirulina could collaborate in prevention of senile dementia. Thus, food products featuring ample amounts of both high-flavanol cocoa powder and spirulina may have considerable potential for health promotion, and merit evaluation in rodent studies and clinical trials.

Vascular-Protective Benefits of Flavanol-Rich Cocoa Powder

A growing number of clinical studies indicate that regular ingestion of flavanol-rich cocoa exerts a range of effects potentially favorable to vascular health – improving endothelial function, reducing elevated blood pressure, increasing insulin sensitivity, and suppressing platelet aggregation.¹⁻¹⁴ There is recent evidence that the epicatechin content of cocoa is primarily responsible for its favorable impact on vascular endothelium, which reflects both an acute and chronic up-regulation of nitric oxide production.^{15, 16} As is well known, physiological levels of nitric oxide support vascular health and efficient tissue perfusion by promoting vasodilation,

opposing inflammation and structural remodeling in the vascular wall, and stabilizing platelets.¹⁷ Other research demonstrates that ingestion of flavanol-rich cocoa protects skin from UV damage and has a positive cosmetic impact on the skin of women, increasing its moisture content.¹⁸

The fact that the Kuna Indians of Panama are virtually immune from hypertension and the typical age-related rise of blood pressure, so long as they live a traditional lifestyle, is likely attributable to their regular heavy intake of flavanol-rich raw cocoa.¹⁰ Moreover, the Kuna appear to be virtually free of stroke¹⁹ – a finding consistent with evidence that cerebrovascular nitric oxide production is a key determinant of stroke risk.²⁰ Although a diet that is nearly pescovegan undoubtedly has a favorable impact on the Kuna's health status, hypertension and stroke tend to be quite common in many Asian groups whose traditional diets are quasi-vegan – even though these groups enjoy considerable protection from coronary heart disease, diabetes, and certain “Western” cancers.²¹⁻²³ It should be emphasized that, unlike all other unacculturated societies that have been found to be free of essential hypertension, the Kunas make ample use of added salt in their diets; thus, cocoa flavanols appear to confer important protection from hypertension and stroke even in the context of a salty diet.

Although the prevalence of senile dementia among the Kuna has not been formally assessed, other Third World cultures in which hypertension and stroke are quite rare are characterized by a near absence of dementia.^{24, 25} This phenomenon may reflect a key role for intermittent or chronic cerebral hypoxia in triggering and sustaining the inflammatory process that mediates Alzheimer's disease. Indeed, many commentators have noted that a number of vascular risk factors are likewise risk factors for Alzheimers, and that many measures which boost endothelial nitric oxide function are linked to decreased risk for this disorder.²⁵⁻³⁵ Recent research has established that hypoxia boosts neuronal expression of BACE1 (a.k.a. beta-secretase), a protease whose activity can be rate-limiting for the production of the amyloid-beta peptides thought to drive the inflammatory process in Alzheimers.^{36, 37} Since amyloid-beta antagonizes endothelium-dependent vasodilation in the cerebral microcirculation by inducing severe endothelial oxidative stress,³⁸⁻⁴² a vicious cycle mechanism may act to sustain and exacerbate local hypoxia (and thus increased BACE1 activity) in regions of the brain where Alzheimers inflammation becomes well established; indeed, endothelial nitric oxide is a mediator of the crucial autoregulatory mechanisms whereby cerebral blood flow is matched to metabolic demand and maintained in the face of reduced central blood pressure.⁴³⁻⁴⁷ A recent clinical study has demonstrated an acute increase of brain perfusion following ingestion of flavanol-rich cocoa.^{48, 49} Moreover, long-term administration of cocoa flavanols to aging rats is associated with preservation of youthful cognitive performance⁵⁰ – raising the possibility that cerebral hypoxia (and/or suboptimal cerebrovascular nitric oxide production) might also play a role in the more modest fall-off in cognitive function associated with healthy aging. These considerations suggest that regular consumption of cocoa flavanols might have important potential for promoting retention of cognitive function as humans age.

Spirulina Has Profound Antioxidant Potential

Another food with potential for superstar status as a health aid is the cyanobacterium spirulina. While spirulina – once harvested as a food by the Aztecs, and still used by Africans living near Lake Chad - has been popular as a supplement in “health food” circles for several decades, its true health-protective merit has only recently been discovered: phycocyanobilin (PCB), the chromophore bound to spirulina’s chief protein, phycocyanin, can function as a potent inhibitor of NADPH oxidase, the enzyme complex that is the chief source of pathological oxidant stress in a wide range of health disorders.^{51, 52} In this regard, it appears to mimic the physiological activity of free bilirubin;⁵³⁻⁵⁶ PCB can be converted within cells to phycocyanorubin, which is nearly identical in structure to bilirubin.⁵⁷ Although the clinical utility of ample intakes of spirulina has so far received little research attention, in numerous rodent studies orally administered spirulina or phycocyanin has shown potent anti-inflammatory, cytoprotective, and anti-atherogenic activities; these effects are most likely attributable to down-regulation of NADPH oxidase activity.^{51, 52, 58, 59} A consideration of the central role of NADPH oxidase over-activity in a range of disorders suggests that ample intakes of spirulina may have preventive and therapeutic potential with respect to many vascular diseases (including atherogenesis, hypertension, and congestive heart failure), cancers, complications of diabetes, and a range of neurodegenerative, fibrotic, or inflammatory disorders.^{51, 52} It should be emphasized that measures which inhibit NADPH oxidase activity could be expected to have a much more profound health impact than antioxidant vitamins or phytochemicals that act merely as oxidant scavengers – the latter, while helping to preserve the structural integrity of membrane lipids or proteins, have little influence on the signal-modulatory activity of hydrogen peroxide, or on the nitric oxide-quenching activity of superoxide. The versatile health protection associated with statin therapy or angiotensin II antagonism – seemingly greater than would be predicted from their hypolipidemic or anti-hypertensive activities – may be largely attributable to their down-regulatory impacts on the NADPH oxidase activity of certain tissues.⁶⁰

Spirulina also is a source of polysaccharide that has immunostimulant activity (reflecting the activation of TLR2 receptors on macrophages),⁶¹⁻⁶³ and is very rich in zeaxanthin, a dietary carotenoid that has been linked to decreased risk for macular degeneration.⁶⁴ A recent open clinical trial reports worthwhile reductions in blood pressure and improvements in blood lipid profile in healthy volunteers receiving 4.5 g spirulina daily; this suggests that spirulina-bound PCB may have good oral bioavailability in humans.⁶⁵ The reduction in LDL cholesterol observed in this study might reflect phycocyanin-mediated inhibition of cholesterol and bile acid (re)absorption, as demonstrated in rats.⁶⁶

Cocoa Can Mask Spirulina's Flavor, While Complementing its Health Benefits

Spirulina's chief drawback as a food is that it has a foul odor, and a flavor that most find unappealing. However, these authors have observed that, when pre-blended with spirulina, cocoa powder can do an excellent job of masking spirulina's odor and flavor. When blended with soy milk, cow's milk, or rice milk, along with an added sweetener, a cocoa-spirulina powder can yield a drink with a rich and creamy chocolate flavor – though some may initially find its dark green color somewhat disconcerting!

Moreover, the antioxidant activity of PCB in the inflamed vasculature – where NADPH oxidase is the chief source of oxidant stress⁶⁷⁻⁶⁹ - could be expected to nicely complement the impact of cocoa-derived epicatechin on nitric oxide bioactivity. As is well known, superoxide antagonizes the bioactivity of nitric oxide by spontaneously reacting with it to generate the dangerous oxidant peroxynitrite. Furthermore, by oxidizing the cofactor tetrahydrobiopterin, oxidant stress (i.e. peroxynitrite) transforms the endothelial nitric oxide synthase into an enzyme that not only is less competent at generating nitric oxide, but that also produces superoxide.^{70, 71} Hence, the protective impact of epicatechin on an inflamed vasculature would be expected to be greater if concurrently administered PCB is employed to quell vascular oxidant stress. And the antioxidant impact of PCB on vascular endothelium would be expected to act in other ways, complementary to but independent of nitric oxide, to minimize endothelial inflammation and thus promote vascular health.^{60, 68, 72, 73}

Oral administration of phycocyanin or of whole spirulina has exerted central neuroprotective effects in rodent studies – an observation which strongly suggests that PCB can transit the blood-brain barrier.⁷⁴⁻⁷⁶ This is of considerable interest in light of evidence that oxidant stress generated by NADPH oxidase in activated microglia and possibly neurons as well plays an important pathogenic role in many common neurodegenerative disorders – including Alzheimers disease.⁷⁶⁻⁸² Indeed, oxidative stress up-regulates transcription of BACE1, an effect mediated by the stress-activated MAP kinases;⁸³⁻⁸⁸ the resulting increase in amyloid-beta production then triggers further oxidative stress via activation of NADPH oxidase, completing a feed-forward loop.⁸⁹⁻⁹⁴ Oxidative stress also promotes transcription of presenilin-1, the catalytic component of the gamma-secretase also required for amyloid-beta production.⁸⁷ Evidently, PCB has the potential to suppress this vicious cycle, and also would likely antagonize the adverse impact of amyloid-beta on cerebrovascular endothelial function, which appears to be mediated by NADPH oxidase activation.⁴² These considerations suggest that cocoa flavanols and PCB could work in tandem to counteract the cerebral hypoxia and oxidative stress that sustain excessive amyloid-beta production and that mediate, at least in part, the neuronal dysfunction and death that characterize Alzheimers.^{77, 78, 89-91, 95-98} Thus, it is conceivable that spirulina could complement the utility of flavanol-rich cocoa in dementia prevention – only by aiding efficient cerebrovascular perfusion, but also by blunting the key contribution of oxidative stress to Alzheimers pathology.

In regard to the UV-protective effect documented for cocoa flavanols, there is suggestive evidence that UV-induced skin damage is mediated by activation of NADPH oxidase in keratinocytes, an effect contingent on concurrent activation of EGF receptors.⁹⁹⁻¹⁰² If this is the case, PCB may be able to complement the utility of cocoa flavanols in serving as an “internal sun screen”.

It is therefore proposed that commercial products combining ample amounts of flavanol-rich cocoa powder and phycocyanin-rich spirulina should be developed, and their effects assessed both in rodent studies and in clinical trials. Consumed regularly, such products may have considerable potential for preventing and treating the wide range of disorders in which excessive oxidative stress plays a pathogenic role – and possibly for preserving youthful cognitive function into ripe old age.

If this proposal proves to have merit, it may ultimately prove feasible to provide the key active components of cocoa powder and spirulina in capsule form. Flavanol-rich cocoa powder extracts are already commercially available, and it seems likely that PCB-enriched spirulina extracts could be developed for use in nutraceuticals.

References

- (1) Taubert D, Berkels R, Roesen R, Klaus W. Chocolate and blood pressure in elderly individuals with isolated systolic hypertension. *JAMA* 2003 August 27;290(8):1029-30.
- (2) Fisher ND, Hughes M, Gerhard-Herman M, Hollenberg NK. Flavanol-rich cocoa induces nitric-oxide-dependent vasodilation in healthy humans. *J Hypertens* 2003 December;21(12):2281-6.
- (3) Engler MB, Engler MM, Chen CY, Malloy MJ, Browne A, Chiu EY, Kwak HK, Milbury P, Paul SM, Blumberg J, Mietus-Snyder ML. Flavonoid-rich dark chocolate improves endothelial function and increases plasma epicatechin concentrations in healthy adults. *J Am Coll Nutr* 2004 June;23(3):197-204.
- (4) Grassi D, Lippi C, Necozione S, Desideri G, Ferri C. Short-term administration of dark chocolate is followed by a significant increase in insulin sensitivity and a decrease in blood pressure in healthy persons. *Am J Clin Nutr* 2005 March;81(3):611-4.
- (5) Vlachopoulos C, Aznaouridis K, Alexopoulos N, Economou E, Andreadou I, Stefanadis C. Effect of dark chocolate on arterial function in healthy individuals. *Am J Hypertens* 2005 June;18(6):785-91.
- (6) Grassi D, Necozione S, Lippi C, Croce G, Valeri L, Pasqualetti P, Desideri G, Blumberg JB, Ferri C. Cocoa reduces blood pressure and insulin resistance and improves endothelium-dependent vasodilation in hypertensives. *Hypertension* 2005 August;46(2):398-405.

- (7) Ding EL, Hutfless SM, Ding X, Girotra S. Chocolate and prevention of cardiovascular disease: a systematic review. *Nutr Metab (Lond)* 2006 January 3;3:2.
- (8) Buijsse B, Feskens EJ, Kok FJ, Kromhout D. Cocoa intake, blood pressure, and cardiovascular mortality: the Zutphen Elderly Study. *Arch Intern Med* 2006 February 27;166(4):411-7.
- (9) Engler MB, Engler MM. The emerging role of flavonoid-rich cocoa and chocolate in cardiovascular health and disease. *Nutr Rev* 2006 March;64(3):109-18.
- (10) Hollenberg K. Vascular action of cocoa flavanols in humans: the roots of the story. *J Cardiovasc Pharmacol* 2006;47 Suppl 2:S99-102.
- (11) Rein D, Paglieroni TG, Wun T, Pearson DA, Schmitz HH, Gosselin R, Keen CL. Cocoa inhibits platelet activation and function. *Am J Clin Nutr* 2000 July;72(1):30-5.
- (12) Innes AJ, Kennedy G, McLaren M, Bancroft AJ, Belch JJ. Dark chocolate inhibits platelet aggregation in healthy volunteers. *Platelets* 2003 August;14(5):325-7.
- (13) Heptinstall S, May J, Fox S, Kwik-Urbe C, Zhao L. Cocoa flavanols and platelet and leukocyte function: recent in vitro and ex vivo studies in healthy adults. *J Cardiovasc Pharmacol* 2006;47 Suppl 2:S197-S205.
- (14) Pearson DA, Holt RR, Rein D, Paglieroni T, Schmitz HH, Keen CL. Flavanols and platelet reactivity. *Clin Dev Immunol* 2005 March;12(1):1-9.
- (15) Schroeter H, Heiss C, Balzer J, Kleinbongard P, Keen CL, Hollenberg NK, Sies H, Kwik-Urbe C, Schmitz HH, Kelm M. (-)-Epicatechin mediates beneficial effects of flavanol-rich cocoa on vascular function in humans. *Proc Natl Acad Sci U S A* 2006 January 24;103(4):1024-9.
- (16) Heiss C, Finis D, Kleinbongard P, Hoffmann A, Rassaf T, Kelm M, Sies H. Sustained increase in flow-mediated dilation after daily intake of high-flavanol cocoa drink over 1 week. *J Cardiovasc Pharmacol* 2007 February;49(2):74-80.
- (17) Cooke JP. The pivotal role of nitric oxide for vascular health. *Can J Cardiol* 2004 August;20 Suppl B:7B-15B.
- (18) Heinrich U, Neukam K, Tronnier H, Sies H, Stahl W. Long-term ingestion of high flavanol cocoa provides photoprotection against UV-induced erythema and improves skin condition in women. *J Nutr* 2006 June;136(6):1565-9.
- (19) Bayard V, Chamorro F, Motta J, Hollenberg NK. Does flavanol intake influence mortality from nitric oxide-dependent processes? Ischemic heart disease, stroke, diabetes mellitus, and cancer in Panama. *Int J Med Sci* 2007;4(1):53-8.
- (20) McCarty MF. Up-regulation of endothelial nitric oxide activity as a central strategy for prevention of ischemic stroke - just say NO to stroke! *Med Hypotheses* 2000 November;55(5):386-403.

- (21) Kuller L, Reisler DM. An explanation for variations in distribution of stroke and arteriosclerotic heart disease among populations and racial groups. *Am J Epidemiol* 1971 January;93(1):1-9.
- (22) Reed DM. The paradox of high risk of stroke in populations with low risk of coronary heart disease. *Am J Epidemiol* 1990 April;131(4):579-88.
- (23) Vartiainen E, Du DJ, Marks JS, Korhonen H, Geng GY, Guo ZY, Koplan JP, Pietinen P, We GL, Williamson D. Mortality, cardiovascular risk factors, and diet in China, Finland, and the United States. *Public Health Rep* 1991 January;106(1):41-6.
- (24) McCarty MF. Up-regulation of endothelial nitric oxide activity as a central strategy for prevention of ischemic stroke - just say NO to stroke! *Med Hypotheses* 2000 November;55(5):386-403.
- (25) McCarty MF. Toward prevention of Alzheimers disease--potential nutraceutical strategies for suppressing the production of amyloid beta peptides. *Med Hypotheses* 2006;67(4):682-97.
- (26) de la Torre JC. Alzheimer's disease is a vasocognopathy: a new term to describe its nature. *Neurol Res* 2004 July;26(5):517-24.
- (27) Luchsinger JA, Reitz C, Honig LS, Tang MX, Shea S, Mayeux R. Aggregation of vascular risk factors and risk of incident Alzheimer disease. *Neurology* 2005 August 23;65(4):545-51.
- (28) Stampfer MJ. Cardiovascular disease and Alzheimer's disease: common links. *J Intern Med* 2006 September;260(3):211-23.
- (29) Rosano C, Newman AB. Cardiovascular disease and risk of Alzheimer's disease. *Neurol Res* 2006 September;28(6):612-20.
- (30) Razay G, Vreugdenhil A, Wilcock G. The metabolic syndrome and Alzheimer disease. *Arch Neurol* 2007 January;64(1):93-6.
- (31) Zhu X, Smith MA, Honda K, Aliev G, Moreira PI, Nunomura A, Casadesus G, Harris PL, Siedlak SL, Perry G. Vascular oxidative stress in Alzheimer disease. *J Neurol Sci* 2007 June 15;257(1-2):240-6.
- (32) Rosendorff C, Beerli MS, Silverman JM. Cardiovascular risk factors for Alzheimer's disease. *Am J Geriatr Cardiol* 2007 May;16(3):143-9.
- (33) Korczyn AD, Vakhapova V. The prevention of the dementia epidemic. *J Neurol Sci* 2007 June 15;257(1-2):2-4.
- (34) Dede DS, Yavuz B, Yavuz BB, Cankurtaran M, Halil M, Ulger Z, Cankurtaran ES, Aytemir K, Kabakci G, Ariogul S. Assessment of endothelial function in Alzheimer's disease: is Alzheimer's disease a vascular disease? *J Am Geriatr Soc* 2007 October;55(10):1613-7.
- (35) Mielke MM, Rosenberg PB, Tschanz J, Cook L, Corcoran C, Hayden KM, Norton M, Rabins PV, Green RC, Welsh-Bohmer KA, Breitner JC, Munger R, Lyketsos CG. Vascular factors predict rate of progression in Alzheimer disease. *Neurology* 2007 November 6;69(19):1850-8.

- (36) Sun X, He G, Qing H, Zhou W, Dobie F, Cai F, Staufenbiel M, Huang LE, Song W. Hypoxia facilitates Alzheimer's disease pathogenesis by up-regulating BACE1 gene expression. *Proc Natl Acad Sci U S A* 2006 December 5;103(49):18727-32.
- (37) Zhang X, Zhou K, Wang R, Cui J, Lipton SA, Liao FF, Xu H, Zhang YW. Hypoxia-inducible factor 1alpha (HIF-1alpha)-mediated hypoxia increases BACE1 expression and beta-amyloid generation. *J Biol Chem* 2007 April 13;282(15):10873-80.
- (38) Stamler JS. Alzheimer's disease. A radical vascular connection. *Nature* 1996 March 14;380(6570):108-11.
- (39) Price JM, Sutton ET, Hellermann A, Thomas T. beta-Amyloid induces cerebrovascular endothelial dysfunction in the rat brain. *Neurol Res* 1997 October;19(5):534-8.
- (40) Thomas T, McLendon C, Sutton ET, Thomas G. beta-Amyloid-induced cerebrovascular endothelial dysfunction. *Ann N Y Acad Sci* 1997 September 26;826:447-51.
- (41) Iadecola C. Cerebrovascular effects of amyloid-beta peptides: mechanisms and implications for Alzheimer's dementia. *Cell Mol Neurobiol* 2003 October;23(4-5):681-9.
- (42) Park L, Anrather J, Zhou P, Frys K, Pitstick R, Younkin S, Carlson GA, Iadecola C. NADPH-oxidase-derived reactive oxygen species mediate the cerebrovascular dysfunction induced by the amyloid beta peptide. *J Neurosci* 2005 February 16;25(7):1769-77.
- (43) Penix LP, Davis W, Subramaniam S. Inhibition of NO synthase increases the severity of kainic acid-induced seizures in rodents. *Epilepsy Res* 1994 July;18(3):177-84.
- (44) Berger C, von KR. Does NO regulate the cerebral blood flow response in hypoxia? *Acta Neurol Scand* 1998 February;97(2):118-25.
- (45) Kajita Y, Takayasu M, Dietrich HH, Dacey RG, Jr. Possible role of nitric oxide in autoregulatory response in rat intracerebral arterioles. *Neurosurgery* 1998 April;42(4):834-41.
- (46) Jones SC, Radinsky CR, Furlan AJ, Chyatte D, Perez-Trepichio AD. Cortical NOS inhibition raises the lower limit of cerebral blood flow-arterial pressure autoregulation. *Am J Physiol* 1999 April;276(4 Pt 2):H1253-H1262.
- (47) White RP, Vallance P, Markus HS. Effect of inhibition of nitric oxide synthase on dynamic cerebral autoregulation in humans. *Clin Sci (Lond)* 2000 December;99(6):555-60.
- (48) Fisher ND, Sorond FA, Hollenberg NK. Cocoa flavanols and brain perfusion. *J Cardiovasc Pharmacol* 2006;47 Suppl 2:S210-S214.
- (49) Francis ST, Head K, Morris PG, MacDonald IA. The effect of flavanol-rich cocoa on the fMRI response to a cognitive task in healthy young people. *J Cardiovasc Pharmacol* 2006;47 Suppl 2:S215-S220.
- (50) Bisson JF, Nejd A, Rozan P, Hidalgo S, Lalonde R, Messaoudi M. Effects of long-term administration of a cocoa polyphenolic extract (Acticoa powder) on cognitive performances in aged rats. *Br J Nutr* 2008 January 8;1-8.

- (51) McCarty MF. "Iatrogenic Gilbert syndrome" - a strategy for reducing vascular and cancer risk by increasing plasma unconjugated bilirubin. *Medical Hypotheses* 2007;accepted for publication.
- (52) McCarty MF. Clinical potential of spirulina as a source of phycocyanobilin. *J Medicinal Food* 2007;in press.
- (53) Jiang F, Roberts SJ, Datla S, Dusting GJ. NO modulates NADPH oxidase function via heme oxygenase-1 in human endothelial cells. *Hypertension* 2006 November;48(5):950-7.
- (54) Matsumoto H, Ishikawa K, Itabe H, Maruyama Y. Carbon monoxide and bilirubin from heme oxygenase-1 suppresses reactive oxygen species generation and plasminogen activator inhibitor-1 induction. *Mol Cell Biochem* 2006 October;291(1-2):21-8.
- (55) Lanone S, Bloc S, Foresti R, Almolki A, Taille C, Callebert J, Conti M, Goven D, Aubier M, Dureuil B, El-Benna J, Motterlini R, Boczkowski J. Bilirubin decreases nos2 expression via inhibition of NAD(P)H oxidase: implications for protection against endotoxic shock in rats. *FASEB J* 2005 November;19(13):1890-2.
- (56) Datla SR, Dusting GJ, Mori TA, Taylor CJ, Croft KD, Jiang F. Induction of Heme Oxygenase-1 In Vivo Suppresses NADPH Oxidase Derived Oxidative Stress. *Hypertension* 2007 August 6.
- (57) Terry MJ, Maines MD, Lagarias JC. Inactivation of phytochrome- and phycobiliprotein-chromophore precursors by rat liver biliverdin reductase. *J Biol Chem* 1993 December 15;268(35):26099-106.
- (58) Romay C, Gonzalez R, Ledon N, Ramirez D, Rimbau V. C-phycocyanin: a biliprotein with antioxidant, anti-inflammatory and neuroprotective effects. *Curr Protein Pept Sci* 2003 June;4(3):207-16.
- (59) Riss J, Decorde K, Sutra T, Delage M, Baccou JC, Jouy N, Brune JP, Oreal H, Cristol JP, Rouanet JM. Phycobiliprotein C-Phycocyanin from *Spirulina platensis* Is Powerfully Responsible for Reducing Oxidative Stress and NADPH Oxidase Expression Induced by an Atherogenic Diet in Hamsters. *J Agric Food Chem* 2007 September 19;55(19):7962-7.
- (60) Guzik TJ, Harrison DG. Vascular NADPH oxidases as drug targets for novel antioxidant strategies. *Drug Discov Today* 2006 June;11(11-12):524-33.
- (61) Balachandran P, Pugh ND, Ma G, Pasco DS. Toll-like receptor 2-dependent activation of monocytes by *Spirulina* polysaccharide and its immune enhancing action in mice. *Int Immunopharmacol* 2006 December 5;6(12):1808-14.
- (62) Pugh N, Ross SA, ElSohly HN, ElSohly MA, Pasco DS. Isolation of three high molecular weight polysaccharide preparations with potent immunostimulatory activity from *Spirulina platensis*, *Aphanizomenon flos-aquae* and *Chlorella pyrenoidosa*. *Planta Med* 2001 November;67(8):737-42.
- (63) Grzanna R, Polotsky A, Phan PV, Pugh N, Pasco D, Frondoza CG. Immolina, a high-molecular-weight polysaccharide fraction of *Spirulina*, enhances chemokine expression in human monocytic THP-1 cells. *J Altern Complement Med* 2006 June;12(5):429-35.

- (64) Ribaya-Mercado JD, Blumberg JB. Lutein and zeaxanthin and their potential roles in disease prevention. *J Am Coll Nutr* 2004 December;23(6 Suppl):567S-87S.
- (65) Torres-Duran PV, Ferreira-Hermosillo A, Juarez-Oropeza MA. Antihyperlipemic and antihypertensive effects of *Spirulina maxima* in an open sample of mexican population: a preliminary report. *Lipids Health Dis* 2007 November 26;6(1):33.
- (66) Nagaoka S, Shimizu K, Kaneko H, Shibayama F, Morikawa K, Kanamaru Y, Otsuka A, Hirahashi T, Kato T. A novel protein C-phycoerythrin plays a crucial role in the hypocholesterolemic action of *Spirulina platensis* concentrate in rats. *J Nutr* 2005 October;135(10):2425-30.
- (67) Brosnan J. Vascular NAD(P)H oxidase as a novel therapeutic target in vascular disease. *Drug News Perspect* 2004 September;17(7):429-34.
- (68) Sorescu D, Szocs K, Griendling KK. NAD(P)H oxidases and their relevance to atherosclerosis. *Trends Cardiovasc Med* 2001 April;11(3-4):124-31.
- (69) Cave AC, Brewer AC, Narayanapanicker A, Ray R, Grieve DJ, Walker S, Shah AM. NADPH oxidases in cardiovascular health and disease. *Antioxid Redox Signal* 2006 May;8(5-6):691-728.
- (70) Milstien S, Katusic Z. Oxidation of tetrahydrobiopterin by peroxynitrite: implications for vascular endothelial function. *Biochem Biophys Res Commun* 1999 October 5;263(3):681-4.
- (71) Kohnen SL, Mouithys-Mickalad AA, by-Dupont GP, Deby CM, Lamy ML, Noels AF. Oxidation of tetrahydrobiopterin by peroxynitrite or oxoferryl species occurs by a radical pathway. *Free Radic Res* 2001 December;35(6):709-21.
- (72) Brandes RP. Role of NADPH oxidases in the control of vascular gene expression. *Antioxid Redox Signal* 2003 December;5(6):803-11.
- (73) Ray R, Shah AM. NADPH oxidase and endothelial cell function. *Clin Sci (Lond)* 2005 September;109(3):217-26.
- (74) Rimbau V, Camins A, Romay C, Gonzalez R, Pallas M. Protective effects of C-phycoerythrin against kainic acid-induced neuronal damage in rat hippocampus. *Neurosci Lett* 1999 December 3;276(2):75-8.
- (75) Chamorro G, Perez-Albiter M, Serrano-Garcia N, Mares-Samano JJ, Rojas P. *Spirulina maxima* pretreatment partially protects against 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine neurotoxicity. *Nutr Neurosci* 2006 October;9(5-6):207-12.
- (76) McCarty MF. Oral phycoerythrin may diminish the pathogenicity of activated microglia in neurodegenerative disorders. *Medical Hypotheses* 2008;in submission.
- (77) Torreilles F, Salman-Tabcheh S, Guerin M, Torreilles J. Neurodegenerative disorders: the role of peroxynitrite. *Brain Res Brain Res Rev* 1999 August;30(2):153-63.

- (78) Mander P, Brown GC. Activation of microglial NADPH oxidase is synergistic with glial iNOS expression in inducing neuronal death: a dual-key mechanism of inflammatory neurodegeneration. *J Neuroinflammation* 2005 September 12;2:20.
- (79) Gao HM, Liu B, Zhang W, Hong JS. Critical role of microglial NADPH oxidase-derived free radicals in the in vitro MPTP model of Parkinson's disease. *FASEB J* 2003 October;17(13):1954-6.
- (80) Qin L, Liu Y, Cooper C, Liu B, Wilson B, Hong JS. Microglia enhance beta-amyloid peptide-induced toxicity in cortical and mesencephalic neurons by producing reactive oxygen species. *J Neurochem* 2002 November;83(4):973-83.
- (81) Shimohama S, Tanino H, Kawakami N, Okamura N, Kodama H, Yamaguchi T, Hayakawa T, Nunomura A, Chiba S, Perry G, Smith MA, Fujimoto S. Activation of NADPH oxidase in Alzheimer's disease brains. *Biochem Biophys Res Commun* 2000 June 24;273(1):5-9.
- (82) Wu DC, Re DB, Nagai M, Ischiropoulos H, Przedborski S. The inflammatory NADPH oxidase enzyme modulates motor neuron degeneration in amyotrophic lateral sclerosis mice. *Proc Natl Acad Sci U S A* 2006 August 8;103(32):12132-7.
- (83) Tamagno E, Bardini P, Obbili A, Vitali A, Borghi R, Zaccheo D, Pronzato MA, Danni O, Smith MA, Perry G, Tabaton M. Oxidative stress increases expression and activity of BACE in NT2 neurons. *Neurobiol Dis* 2002 August;10(3):279-88.
- (84) Tong Y, Zhou W, Fung V, Christensen MA, Qing H, Sun X, Song W. Oxidative stress potentiates BACE1 gene expression and Abeta generation. *J Neural Transm* 2005 March;112(3):455-69.
- (85) Tamagno E, Parola M, Bardini P, Piccini A, Borghi R, Guglielmotto M, Santoro G, Davit A, Danni O, Smith MA, Perry G, Tabaton M. Beta-site APP cleaving enzyme up-regulation induced by 4-hydroxynonenal is mediated by stress-activated protein kinases pathways. *J Neurochem* 2005 February;92(3):628-36.
- (86) Coma M, Guix FX, Ill-Raga G, Uribealago I, Alameda F, Valverde MA, Munoz FJ. Oxidative stress triggers the amyloidogenic pathway in human vascular smooth muscle cells. *Neurobiol Aging* 2007 February 14.
- (87) Tamagno E, Guglielmotto M, Aragno M, Borghi R, Autelli R, Giliberto L, Muraca G, Danni O, Zhu X, Smith MA, Perry G, Jo DG, Mattson MP, Tabaton M. Oxidative stress activates a positive feedback between the gamma- and beta-secretase cleavages of the beta-amyloid precursor protein. *J Neurochem* 2008 February;104(3):683-95.
- (88) Tamagno E, Guglielmotto M, Giliberto L, Vitali A, Borghi R, Autelli R, Danni O, Tabaton M. JNK and ERK1/2 pathways have a dual opposite effect on the expression of BACE1. *Neurobiol Aging* 2008 February 5.
- (89) Shimohama S, Tanino H, Kawakami N, Okamura N, Kodama H, Yamaguchi T, Hayakawa T, Nunomura A, Chiba S, Perry G, Smith MA, Fujimoto S. Activation of NADPH oxidase in Alzheimer's disease brains. *Biochem Biophys Res Commun* 2000 June 24;273(1):5-9.

- (90) Qin L, Liu Y, Cooper C, Liu B, Wilson B, Hong JS. Microglia enhance beta-amyloid peptide-induced toxicity in cortical and mesencephalic neurons by producing reactive oxygen species. *J Neurochem* 2002 November;83(4):973-83.
- (91) Jana A, Pahan K. Fibrillar amyloid-beta peptides kill human primary neurons via NADPH oxidase-mediated activation of neutral sphingomyelinase. Implications for Alzheimer's disease. *J Biol Chem* 2004 December 3;279(49):51451-9.
- (92) Park L, Anrather J, Zhou P, Frys K, Pitstick R, Younkin S, Carlson GA, Iadecola C. NADPH-oxidase-derived reactive oxygen species mediate the cerebrovascular dysfunction induced by the amyloid beta peptide. *J Neurosci* 2005 February 16;25(7):1769-77.
- (93) Wilkinson B, Koenigsnecht-Talboo J, Grommes C, Lee CY, Landreth G. Fibrillar beta-amyloid-stimulated intracellular signaling cascades require Vav for induction of respiratory burst and phagocytosis in monocytes and microglia. *J Biol Chem* 2006 July 28;281(30):20842-50.
- (94) Abramov AY, Duchen MR. The role of an astrocytic NADPH oxidase in the neurotoxicity of amyloid beta peptides. *Philos Trans R Soc Lond B Biol Sci* 2005 December 29;360(1464):2309-14.
- (95) de la Torre JC. Alzheimer's disease is a vasocognopathy: a new term to describe its nature. *Neurol Res* 2004 July;26(5):517-24.
- (96) Zhang YJ, Xu YF, Liu YH, Yin J, Li HL, Wang Q, Wang JZ. Peroxynitrite induces Alzheimer-like tau modifications and accumulation in rat brain and its underlying mechanisms. *FASEB J* 2006 July;20(9):1431-42.
- (97) Malinski T. Nitric oxide and nitroxidative stress in Alzheimer's disease. *J Alzheimers Dis* 2007 May;11(2):207-18.
- (98) Peers C, Pearson HA, Boyle JP. Hypoxia and Alzheimer's disease. *Essays Biochem* 2007;43:153-64.
- (99) Chamulitrat W, Stremmel W, Kawahara T, Rokutan K, Fujii H, Wingler K, Schmidt HH, Schmidt R. A constitutive NADPH oxidase-like system containing gp91phox homologs in human keratinocytes. *J Invest Dermatol* 2004 April;122(4):1000-9.
- (100) Wang H, Kochevar IE. Involvement of UVB-induced reactive oxygen species in TGF-beta biosynthesis and activation in keratinocytes. *Free Radic Biol Med* 2005 April 1;38(7):890-7.
- (101) Beak SM, Lee YS, Kim JA. NADPH oxidase and cyclooxygenase mediate the ultraviolet B-induced generation of reactive oxygen species and activation of nuclear factor-kappaB in HaCaT human keratinocytes. *Biochimie* 2004 July;86(7):425-9.
- (102) Valencia A, Kochevar IE. Nox1-based NADPH oxidase is the major source of UVA-induced reactive oxygen species in human keratinocytes. *J Invest Dermatol* 2008 January;128(1):214-22.